



### Risankizumab (Skyrizi) Order Set for Active Crohn's disease (CD):

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_

Height: \_\_\_\_\_ Weight: \_\_\_\_\_ (kg) Allergies: \_\_\_\_\_

#### Diagnosis:

\_\_\_\_ K50. \_\_\_\_ Crohn's Disease \_\_\_\_ Other (ICD-10 Code: \_\_\_\_\_)

#### Orders for Outpatient Infusion:

- **Fax order to 337-430-6976 once medical prior authorization obtained**
- **Assign as Outpatient**
- **Criteria for Administration:**
  - A negative TB skin test or other appropriate documentation of TB status must be faxed to 430-6976 prior to scheduling of appointment for patient.
- **Nursing:** Confirm TB and hepatitis B status (or has received hepatitis B vaccination). Assess patient for active infection prior to initiation of therapy: notify MD if present.
- **Labs:** CMP at baseline and prior to every IV infusion
- **Premedications – Give 15 minutes prior to infusion:**

____ Acetaminophen 650 mg PO x 1 dose	____ methylPREDNISolone 125 mg IV x 1 dose
____ diphenhydrAMINE 25 mg PO x 1 dose	____ diphenhydrAMINE 25 mg IV x 1 dose
____ diphenhydrAMINE 50 mg PO x 1 dose	____ diphenhydrAMINE 50 mg IV x 1 dose
____ Other: _____	
- **Risankizumab Induction Dosing**
  - Risankizumab (Skyrizi) (J3590) 600 mg in D5W 100 ml IV over 60 minutes at weeks 0, 4 and 8 weeks.
- **Severe Reactions:** Stop infusion, initiate anaphylaxis protocol and notify MD.
- **IV Line Care:**
  - Normal Saline 10 ml IV flush after each use
  - For implanted ports: Heparin 100 units/ml 5 ml IV flush after each use or prior to deaccessing
- Discharge when infusion complete

Physician Signature: \_\_\_\_\_ Date/Time: \_\_\_\_\_

#### Orders for Specialty Pharmacy – FOR specialty pharmacy use ONLY:

- **Fax order to 337-494-6536 (pharmacy will get prior authorization for pharmacy benefits)**

Risankizumab (Skyrizi) On-Body Injector with prefilled cartridge - Choose one:

(use lowest *effective* dose to maintain therapeutic response)

\_\_\_\_ Inject 180 mg SUBQ into thigh or abdomen on week 12 and then every 8 weeks

\_\_\_\_ Inject 360 mg SUBQ into thigh or abdomen on week 12 and then every 8 weeks

Quantity: #1 Refills: \_\_\_\_\_

\_\_\_\_ Generic substitution permitted \_\_\_\_ Dispense as Written

Physician Signature: \_\_\_\_\_ Date/Time: \_\_\_\_\_



Patient: «Full\_Name»; DOB: «Birth\_Date»

Physician: «Attending\_Physician\_Last\_Name», «Attending\_Physician\_First\_Name» «Attending\_Physician\_Middle\_Init»

Visit ID: «Visit\_ID»