



PO_SYSTEMIC LUPUS ERYTHMATOSUS (SLE) ORDER SET Last Revised: 01/10/2023

	Erythematosus (S			DOB:
Height:	Weight:	(kg)	Allergies:	DOB:
Assign as Outpa				
Diagnosis:				
M32.14 Lupu				_M32.8 Systemic Lupus Erythematosus
Other (ICD-1	10 Code):			
Labs: To be done	e per MD office as C	Outpatient prio	r to admittand	ce to Infusion Center.
Nursing: Assess	patient for active ir	nfection prior t	o initiation of	therapy: notify MD if present.
Belimumal	b (Benlysta)			
Pre-Medic				
	etaminophen 650 m	•	•	
-	henhydrAMINE 25	mg PO x 1 do	se prior to inf	fusion
	tial Dosing:	. NO 050 11		
mg				r every 2 weeks x 3 doses, then belimumab 10 to the nearest vial size if no greater than 10%
	sequent Dosing:			
		in NS 250 ml I	IV over 1 hou	ir every 4 weeks. Doses may be rounded to the
	arest vial size if no			
Minor rea	ctions: Slow the in	fusion rate or	temporarily in	nterrupt the infusion.
Severe Re	eactions: Stop infu	sion, initiate a	anaphylaxis p	rotocol and notify MD.
Anifroluma	ab-fnia (Saphnelo)			
	cations:			
				30 minutes using a 0.2 micron filter every 4 week
	sion set with NS 25			rotocal and notify MD
• Severe Re	sactions: Stop into	ision, initiate a	апарпувахіѕ р	rotocol and notify MD.
IV Line Care:				
	aline 10 ml IV flush a nted ports: Heparin			after each use or prior to deaccessing
Discharge when i	nfusion complete			
New MD order re	equired every 6 mor	nths unless de	fined in origin	nal order
Physician Signatu	iro.			Date/Time:
i nysician signatt	п с.			Date/Time:



